



## SUCRASE (MALTASE) + $\beta$ -GALACTOSIDASE (Lot 191106)

E-SUCRBG

05/20

**Sucrase (170 U),  $\alpha$ -glucosidase (2600 U) and  $\beta$ -galactosidase (3000 U); freeze dried.**

For use in the removal of sucrose, maltose and lactose in dietary fiber determinations. In the [Integrated Total Dietary Fiber procedure](#), in HPLC analysis of non-digestible oligosaccharides using the Waters Sugar-Pak column, the fructosyl-trisaccharide  $\beta$ -D Fruf (2 $\rightarrow$ 1)- $\beta$ -D-Fruf-(2 $\rightarrow$ 1)- $\beta$ -D-Fruf, chromatographs at a similar point to the disaccharides, sucrose, maltose and lactose. Accurate determination of this trisaccharide requires the hydrolysis of these disaccharides. This can be achieved using this enzyme mixture.

### PROPERTIES

**1. ELECTROPHORETIC PURITY:**

This is a mixture of sucrase (maltase; from yeast):

- Single major band on SDS-gel electrophoresis (57,750)  
plus  $\beta$ -galactosidase (from *A. niger*)
- Single band on isoelectric focusing

**2. ACTIVITY:**

This enzyme mixture gives complete hydrolysis of sucrose, maltose and lactose under the defined assay conditions, with no hydrolysis of the trisaccharide,  $\beta$ -D-Fruf-(2 $\rightarrow$ 1)- $\beta$ -D-Fruf-(2 $\rightarrow$ 1)- $\beta$ -D-Fruf.

**3. STORAGE CONDITIONS:**

The enzyme is supplied as a lyophilised powder and should be stored below -10°C. On dissolution in buffer or water, the enzyme should be stored in the frozen state. It is recommended that all buffers used for dilution contain BSA (0.5 mg/mL).

**4. PREPARATION OF ENZYME FOR USE:**

Dissolve the contents of one vial in 6 mL of 5 mM sodium acetate buffer (pH 5.0). Transfer aliquots of approx. 2 mL to polypropylene tubes and store below -10°C between use. Can be thawed and re-frozen several times.

## INCUBATION CONDITIONS:

To 1 mL of sugar mixture obtained in the **Integrated Total Dietary Fiber procedure [Step 1(b)]** containing up to 5 mg/mL of sucrose, maltose and/or lactose and fructo-triose,

**add:**

0.1 mL of sucrase/ $\beta$ -galactosidase enzyme mixture, and incubate at 40°C, for 60 min.

Terminate the reaction by incubating the tube at 100°C for 2 min and centrifuge the suspension in a Microfuge at 12,000 rpm for 5 min.

## SAMPLE PREPARATION AND HPLC:

Analyse the supernatant solution by HPLC using a Waters Sugar-Pak<sup>R</sup> column as described in the Megazyme kit data booklet for the Integrated Total Dietary Fiber method (**K-INTDF**). Calculate the amount of fructo-triose by reference to the D-sorbitol internal standard. This amount should then be added to the determined amount of non-digestible oligosaccharides [NDO; low molecular weight soluble dietary fiber; Soluble Dietary Fiber soluble in 78% aqueous ethanol (SDFS)].

### Hydrolysis of Sucrose, Maltose and Lactose with Sucrase + $\beta$ -Galactosidase

