


Requirements:

- Cellulase Assay Kit (**K-CellG5-4V / K-CellG5-2V**) (provides ~ 480 assays or 240 assays).
- K-CellG5** ChemWell® 2910 assay file.
- Use in association with the Cellulase Assay Kit (CELLG5 Method) (**K-CellG5-4V or K-CellG5-2V**) product data booklet.

Use:

A specific and sensitive colourimetric method for the determination of cellulase (*endo*-1,4- β -glucanase).

Assay Principle:



Procedure:

Prepare the assay reagents and use with the **K-CellG5** ChemWell® 2910 assay file.

Cellulase Assay Kit Components:

- Bottle 1:** 4,6-O-(3-Ketobutylidene)-4-nitrophenyl- β -D-cellopentaoside (BPNPG5) in 10% DMSO/H₂O (3 mL) plus sodium azide (0.02% w/v). Stable for > 4 years at -20°C.
- Bottle 2:** Thermostable β -glucosidase suspension (0.50 mL, 600 U/mL). Stable for > 4 years at 4°C.
- Bottle 3:** *Trichoderma* cellulase standard solution (5 mL; ~ 3 U/mL; actual value stated on the vial label) in 50% aqueous glycerol plus sodium azide (0.02% w/v). Stable for > 4 years at -20°C.

Preparation of Kit Components:

- 1 & 2.** The CellG5 reagent solution is prepared by transferring 100 μ L of the suspension in bottle 2 (swirl bottle 2 to mix contents before use) to bottle 1 and mix well. The CellG5 reagent solution should only be prepared prior to use. Stable for ~ 4 weeks at 4°C or stable for ~ 5 days at room temperature.
- 3.** With a positive displacement pipette, dispense 0.5 mL of the contents of bottle 3 to 9.5 mL of **Buffer B** and mix well. Once diluted, the standard is stable for ~ 4 days at 4°C or ~ 6 months at -20°C.

Preparation of Reagents Not Supplied:

Stopping Reagent

(500 mM Sodium Carbonate, pH 11.0)

Dissolve 53 g of sodium carbonate (anhydrous) in 1 L of distilled water and adjust the pH to approx. 11.0. Store in a sealed bottle. Stable for ~ 3 months at room temperature.

(A) Concentrated Acetate Buffer

(Sodium acetate buffer, 1 M, pH 4.5)

Add 60 g of glacial acetic acid (1.05 g/mL) to 800 mL of distilled water. Adjust the pH of this solution to 4.5 by the addition of 5 M (20 g/100 mL) NaOH solution. Adjust the volume to 1 L. Stable for > 2 years at room temperature.

(B) Acetate Extraction/Dilution Buffer

(Sodium acetate buffer, 100 mM, pH 4.5 containing 0.02% w/v sodium azide)

Add 100 mL of concentrated acetate buffer (A) to 850 mL of distilled water. Adjust the pH to 4.5 by dropwise addition of 2 M HCl or 2 M NaOH and adjust the volume to 1 L. Add 0.2 g of sodium azide and dissolve. Stable for > 1 year at 4°C.

(C) Concentrated Phosphate Buffer

(Sodium phosphate buffer, 0.5 M, pH 6.0)

Add 156 g of sodium dihydrogen orthophosphate (NaH₂PO₄·2H₂O) to 1.5 L of distilled water. Adjust the pH to 6.0 with 4 M NaOH and adjust the volume to 2 L. Stable for > 1 year at 4°C.

(D) Phosphate Extraction/Dilution Buffer

(Sodium phosphate buffer, 100 mM, pH 6.5 containing 0.02% sodium azide)

Add 200 mL of concentrated phosphate buffer (C) to 750 mL of distilled water. Adjust the pH to 6.5 with 1 M HCl or 1 M NaOH and adjust the volume to 1 L. Add 0.2 g of sodium azide and dissolve. Stable for > 1 year at 4°C.

Extraction and Assay of Milled Grain Samples:

- Add 1.0 mL of **liquid enzyme** preparation to 49 mL of Extraction/Dilution **Buffer B** (pH 4.5) or **Buffer D** (pH 6.0) using a positive displacement dispenser (these solutions can be very viscous) and mix thoroughly. **This is termed the Original Extract.**

Alternatively:

- Add 1.0 g of **powder enzyme** sample to 50 mL of **Buffer B** (pH 4.5) or **Buffer D** (pH 6.0) and gently stir the slurry over a period of approx. 15 min or until the sample is completely dispersed or dissolved. Clarify this solution by centrifugation (1,000 x g, 10 min) or by filtration through Whatman No. 1 (9 cm) filter circles. **This is termed the Original Extract.**
- If further dilutions are required, add 1.0 mL of the **Original Extract** to 9.0 mL of Extraction/Dilution **Buffer B** (pH 4.5) or **Buffer D** (pH 6.0) (10-fold dilution) and mix thoroughly. This process of dilution should be repeated until a suitable concentration of cellulase for assay is achieved.
- Perform the assay using the **K-CellG5** ChemWell® 2910 assay file.





CELLULASE (K-CellG5) Procedure for ChemWell® 2910 Auto-Analyser

Assay Parameters:

Assay volumes: CellG5 Reagent: 0.025 mL
Sample: 0.025 mL
Stopping Reagent: 0.250 mL

Reaction time: 10 min at 37°C

Wavelength: 405 nm

Assay type: Stopped reaction

Reaction direction: Increase

Calculation of Activity:

One Unit of activity is defined as the amount of enzyme, in the presence of excess thermostable β -glucosidase, required to release one micromole of 4-nitrophenol from CellG5 in one minute under the defined assay conditions, and is termed an **CellG5 Unit**.

CellG5 Units/mL or g of original enzyme preparation:

$$= \frac{\Delta E_{405}}{\text{Incubation Time}} \times \frac{\text{Total Volume in Cell}}{\text{Aliquot Assayed}} \times \frac{1}{\epsilon_{\text{mM}}} \times \frac{\text{Extraction Vol.}}{\text{Sample Weight}} \times D$$

where:

ΔE_{405} = Absorbance (reaction) - Absorbance (blank)

Incubation Time = 10 min

Total Volume in Cell = 0.3 mL

Aliquot Assayed = 0.025 mL

ϵ_{mM} of *p*-nitrophenol (at 405 nm) in 500 mM sodium carbonate, pH 11 = 14.652

D = Dilution of the extract including the original extraction (e.g. 50 mL per 1 mL or 1 g of cellulase)

Thus:

CellG5 Units/mL or g:

$$= \frac{\Delta E_{405}}{10} \times \frac{0.300}{0.025} \times \frac{1}{14.652} \times D$$
$$= \Delta E_{405} \times 0.0819 \times D$$

NOTE:

The absorption coefficient (ϵ_{mM}) of 14.652 was experimentally determined under the conditions of the automated cellulase assay (CellG5 method) using a ChemWell® 2910 auto-analyser.

