



α -D-GLUCURONIDASE from *Geobacillus stearothermophilus* (Lot 181101a)

Recombinant - Thermostable

E-AGUBS

12/18

Fusion protein of α -D-glucuronidase
(EC 3.2.1.139) alpha-D-glucosiduronate glucuronohydrolase
CAZy Family: GH67
CAS: 37259-81-7

PROPERTIES

1. ELECTROPHORETIC PURITY:

- Single band on SDS-gel electrophoresis (MW ~ 93,200)
- Broad diffuse band on isoelectric focusing (pI ~ 5.4)

2. SPECIFIC ACTIVITY AND LEVEL OF OTHER ACTIVITIES:

40 U/mg protein (on Aldotriouronic acid) at pH 7.0 and 70°C

10 U/mg protein (on Aldotriouronic acid) at pH 7.0 and 40°C

One Unit of α -D-glucuronidase activity is defined as the amount of enzyme required to release one μ mole of α -D-glucuronic acid per minute from aldotriouronic acid in MOPS buffer (100 mM) pH 7.0 and 40 or 70°C. The assay was performed using the α -D-Glucuronidase Assay Kit from Megazyme (**Megazyme catalogue code: K-AGLUA**).

3. SPECIFICITY:

Hydrolysis of the α -1,2 glycosidic bond between D-glucuronic acid or its ether 4-O-methyl-D-glucuronic acid and D-xylose residues of xylo-oligosaccharides (aldo-uronic acids) from xylan.

4. PHYSICOCHEMICAL PROPERTIES:


pH Optima:	7.0
pH Stability:	6.0 - 9.0 (> 75% control activity after 24 hours at 4°C)
Temperature Optima:	70°C (10 min. reaction)
Temperature Stability:	up to 70°C (> 90% control activity after 15 min.)

5. STORAGE CONDITIONS:


The enzyme is supplied in 50% glycerol containing 0.02% (w/v) sodium azide and should be stored below -10°C. For assay, this enzyme should be diluted in MOPS buffer (100 mM), pH 7.0 containing 0.5 mg/mL BSA.

6. EXPERIMENTAL DATA


pH Optima



pH Stability



Thermal Optima



Thermal Stability

