



β-GLUCOSIDASE from *Thermotoga maritima* (Lot 151102a)

Recombinant - Thermostable

E-BGOSTM

(EC 3.2.1.21) beta-D-glucoside glucohydrolase

CAZy Family: GH1

11/18

PROPERTIES

1. ELECTROPHORETIC PURITY:

- Single band on SDS-gel electrophoresis (MW ~ 53,700)
- One major band on isoelectric focusing (pI ~ 6.0)

2. SPECIFIC ACTIVITY:

50 U/mg protein (on *p*-NP-β-D-Glucopyranoside) at pH 6.5 and 40°C.

One Unit of β-glucosidase activity is defined as the amount of enzyme required to release one μmole of *p*-nitrophenol (*p*-NP) per minute from *p*-nitrophenyl-β-D-glucopyranoside (10 mM) in sodium maleate buffer (50 mM), pH 6.5 at 40°C.

3. OTHER ACTIVITIES (as a percentage of β-glucosidase activity):

Enzyme Measured	Substrate	Activity, %
β-Glucosidase	<i>p</i> -NP-β-D-Glucopyranoside	100
β-Glucosidase	Cellobiose	~ 17
β-Galactosidase	<i>p</i> -NP-β-D-Galactopyranoside	~ 100
α-Amylase	Ceralpha Reagent	< 0.0001
Amyloglucosidase	Starch	< 0.0002
α-Glucosidase	<i>p</i> -NP-α-D-Glucopyranoside	< 0.001
endo-1,4-β-Glucanase	Cellzyme C Tablets (Megazyme)	< 0.0001

Action on polysaccharide and *p*-nitrophenyl substrates was determined at final concentrations of 10 mg/mL and 10 mM, respectively, in sodium maleate buffer (100 mM), pH 6.5 at 40°C.

4. RELATIVE RATES OF HYDROLYSIS OF SUBSTRATES:

Substrate	%
Cellobiose	100
Laminaribiose	~ 109
Laminaritriose	~ 100
Laminaritetraose	~ 59
Laminaripentaose	~ 29
Laminarihexaose	~ 5.0
Gentiobiose	~ 5.9
Sophorose	~ 98
1,4-β-D-Glucosyl-D-mannose	~ 9.9
<i>p</i> -Nitrophenyl β-D-glucopyranoside	~ 566
<i>p</i> -Nitrophenyl β-D-xylanopyranoside	~ 1.8
<i>p</i> -Nitrophenyl α-D-glucopyranoside	< 0.001

5. PHYSICOCHEMICAL PROPERTIES:

pH Optima: 7.0 (at 40°C) Temperature Optima: 90°C (30 min at pH 6.5)
 pH Stability: 4.0-9.0 (at 40°C for 30 min) Temperature Stability: Unstable above 90°C

6. STORAGE CONDITIONS

The enzyme is supplied as an ammonium sulphate suspension in 0.02% (w/v) sodium azide and should be stored at 4°C. For assay, this enzyme should be diluted in sodium maleate buffer (50 mM), pH 6.5 containing 0.5 mg/mL BSA. **Swirl to mix the enzyme immediately prior to use.**