



CELLULASE (*endo*- β -GLUCANASE) from *T. maritima* (Lot 100201d)

Recombinant - Thermostable

E-CELTTM

04/20

EC: 3.2.1.4

Synonyms: cellulase; 4-beta-D-glucan 4-glucanohydrolase

CAZy: GH5

CAS: 9012-54-8

PROPERTIES

1. ELECTROPHORETIC PURITY:

- Single band on SDS-gel electrophoresis (MW ~ 38,200)
- Single major band on isoelectric focusing (pI ~ 6.2)

2. SPECIFIC ACTIVITY:

245 U/mg protein (on CM-Cellulose 4M) at pH 6.0 and 80°C

53 U/mg protein (on CM-Cellulose 4M) at pH 6.0 and 40°C

One Unit of cellulase activity is defined as the amount of enzyme required to release one μ mole of glucose reducing-sugar equivalents per minute from CM-Cellulose 4M (10 mg/mL) in sodium phosphate buffer (100 mM) pH 6.0.

3. RELATIVE RATES OF HYDROLYSIS OF SUBSTRATES:

Substrate	%
CM-Cellulose 4M	100
Carob Galactomannan (low viscosity)	~ 2.5
Wheat Arabinoxylan	< 0.009
pNP- β -D-glucoside	< 0.007

Action on polysaccharide and pNP substrates was determined at final substrate concentrations of 5 mg/mL and 5 mM, respectively, in sodium phosphate buffer (100 mM), pH 6.0 at 40°C.

4. PHYSICOCHEMICAL PROPERTIES:

- pH Optima: 6.0
pH Stability: 3.0-9.0 (> 75% control activity after 24 h at 4°C)
Temperature Optima: 80°C (10 min reaction)
Temperature Stability: up to 80°C (> 90% control activity after 15 min)

5. STORAGE CONDITIONS:

The enzyme is supplied as an ammonium sulphate suspension in 0.02% (w/v) sodium azide and should be stored at 4°C. For assay, this enzyme should be diluted in sodium phosphate buffer (100 mM), pH 6.0 containing 1 mg/mL BSA. **Swirl to mix the enzyme immediately prior to use.**